

HEAT EXHAUSTION/HEAT STROKE

Strenuous activity in the heat may cause heat-related illness. Symptoms may include:

- Red, hot, dry skin.
- Weakness and fatigue.
- Cool, clammy hands.
- Vomiting.
- Loss of consciousness.
- Profuse sweating.
- Headache.
- Nausea.
- Confusion.
- Muscle cramping.

Wear disposable gloves when exposed to body fluids.

Quickly remove the student from heat to a cooler, shaded place.

- Have the student lie down.
- Elevate legs 8-12 inches.

Is the student:

- Unconscious or losing consciousness?
- Hot, dry, have red skin?
- Vomiting?
- Confused?

NO

YES

CALL EMS/911

- Give cool, clear fluids such as water, or commercial electrolyte drink frequently in small amounts if person is fully awake and alert.
- Sponge student with cool wet cloths on head, face, and trunk, change the cloths frequently.
- Fan student.
- Loosen clothing.
- Remove any additional layers of clothing

- Put the student on his/her side to protect the airway.
- Look, listen and feel for breathing.
- If student stops breathing, start CPR.

Contact responsible school authority & parent/guardian.

- Remove any additional layers of clothing.
- Cool rapidly by completely wetting clothing with cool water and fan student.
- **DO NOT USE ICE WATER.**
- Place ice packs on neck, armpits, and groin.
- Give nothing by mouth.

Document care provided.

If unable to reach parent/guardian have student rest with adult supervision and continue to provide clear fluids.